

UDC: 633.1+655.852.53

**EFFECT OF STIMULATORS AND SILICON COMPLEX FERTILIZERS ON THE  
PLACEMENT OF FIRST POD OF SOYBEAN VARIETIES**

**Kamalova Nuriya Shamilevna** - PhD student of Tashkent State Agrarian University,  
Uzbekistan

**Abstract.** This article presents data on the effect of separate and combined application of stimulants and silicon complex fertilizers on soybean varieties on their height of the first pod. It was found that stimulants and silicon complex fertilizers had a good effect on the number of height of the first pod, and in both varieties, the height of the first pod was higher when using Maltamin from the stimulants, when comparing stimulants with silicon complex fertilizer, and when using stimulants with silicon complex, silicon complex fertilizer + Maltamin was taken highest placement of the first pod.

**Keywords:** soybean, stimulants, Uzbiogumin, Regoplant, Maltamin, Sila silicon, height of the first pod, Nafis, Vilana.

**INTRODUCTION**

Soybean plant is a very important plant in the world today. It is grown in more than 60 areas of the world. Soy is the leading crop among legumes. Today, when there is a protein shortage all over the world, the protein content of soybeans, the presence of all the amino acids useful for humans in the protein content, increases the nutritional value of soybeans even more. It is important to note that the advantage of soybeans is that they can be compared with a number of food products in terms of the richness of lysine, methionine, arginine, leucine and other essential amino acids. In many countries where soybeans are grown, this crop is the only source of protein, and it also provides livestock with nutritious food and increases its productivity. Soybean accounts for 40% of the world's gross vegetable oil production [2].

In addition to the adopted cultivation technology, the application and study of stimulants that activate the plant's more active growth, development and crop formation remains a very relevant issue today.

this indicator can be increased under the influence of various stimulants and growth regulators, fertilizers, substances with various active effects [1].

Currently, various complexes are being created at the industrial level, including mineral, micro-fertilizers, growth regulators, stimulants, and seed adhesives. This set of complex substances is used in seed treatment. Seeds are saved when treated in this way. The use of a complex of growth regulators should ensure not only the productivity of the plant, but also its safety [7].

The use of plant growth regulators in agriculture began in the 30s of the last century in the Pieceed States. The first widely used synthetic hormone was ethylene. Since then, synthetic substances imitating natural growth hormones have become an integral part of modern agriculture [4].

The development of the technology of using biostimulants for leguminous crops, which regulate growth and increase immunity, is of the most urgent importance [6].

Achievements achieved in the experiments of "Aksayskaya Niva" LLC, Rostov Region, Aksay District. Treatment of soybean seeds with Gumimax drug had a positive effect on the reduction of wilting in soybeans, plant grains and grain yield. The results of the experiment show that the drug "Gumimax" increases the plant's resistance to existing discomforts. Note: pre-sowing seeds simultaneously with rhizobium treatment of the plant allows to increase the productivity of soybean grain up to 0.3–0.4 tons [3].

Growth regulators have the ability to positively affect the yield and quality of soybean seeds. They increase the resistance of the plant to water, temperature and other adverse conditions. In order to achieve the maximum effect, seeds are obtained by step-by-step processing [8].

Ran O.P., Selikhova O.A., Tikhonchuk P.V. (2009) noted that when soybeans are grown in arid regions, they are not irrigated during the rainy season. In some regions, due to the effect of drought, the harvest is reduced, a lot of damage is seen. Damage from drought depends on the duration of the drought period, the period of plant development, evaporation and soil conditions. Water deficit (drying of the soil) is expressed by the closing of the leaf apices, the reduction of transpiration and photosynthesis [5].

Kh.N. Atabaeva., F.B. Namozov., A.A. Kurbanov and S.Sh. Khayrullaev in their experiments conducted in 2018-2020, when they applied micronutrients to the soybean crop, micronutrients affected the height of the soybean stem, leaf, root development, nodule formation, grain quality and productivity, and provided a high yield [10].

According to R.Jo'raeva., J.Toshpol'atov., A.Iminov., Kh.Bozorov and L.Zaynitdinova, S.Khatamov and S. Sh.Khayrullaev, in their experiments conducted in 2015-2017, soybean plant mineral fertilizers and belonging to the rhizobium group it was observed that the yield increased by 12.6-12.8 c/ha when exposed to azotobacteria strains compared to the control variant [11, 14].

According to Khayrullayev Sardor Shamsiddin ugli (2021), the application of micronutrients in the suspension method 2 times during the application period of soybean varieties in the conditions of meadow-swamp soils provides an increase in grain quality [16]. According to data of Atabayeva Khalima Nazarovna, Khayrullaev Sardor Shamsiddin o'g'li, and Usmonova Shohista Usmon qizi (2020), sulfur has a positive effect on the branching of soybean varieties on the background of mineral fertilizers, and in 2018 the number of branches in the variety "Orzu" increased by 0.8-1.3 compared to the control option due to the micro element sulfur. In the "Nafis" variety, this figure was 0.3-0.4, and good results were obtained from medium and high sulfur standards. In 2019, these indicators increased by 0.3-0.7 in the variants of sulfur compared to the control in the "Orzu" variety, increased by 0.1-0.3 in the "Nafis" variety, and good results were obtained from the medium and high standards of sulfur [13]. According to Iminov Abduvali Abdumannobovich, Khayrullayev Sardor Shamsiddin ugli, et al, Nitragine treatment of soybean and mung bean seeds before sowing had a positive effect on seed germination under both laboratory and field conditions, the germination rate of seeds in the laboratory under the conditions of cotton cultivation in the following year under the background of non-treatment by nitragine before sowing the seeds of soybean and mung bean crops grown as a secondary crop after winter wheat was 0.3-1.3%, and field fertility was 0.2-0.8% higher. Also, it was

found that the use of phosphorus and potassium fertilizers in soybean and mung bean crops grown as a secondary crop was 0.6-1.0% higher in the laboratory, and 0.6-0.7% higher in the field than in the control options without mineral fertilizers in studies [12]. According to Umarova Nigora Sadriddinovna, Bo'riboev Bekzod Yetmish ugli, Khayrullayev Sardor Shamsiddin ugli, Usmonova Shokhista Usmon kizi, & Turdaliyeva Shokhista Tulkinjon kizi, the demand of the soybean plant for mineral fertilizers, it was observed that when NPK and liquid fertilizer were used together, all the biometric parameters and yields of the plant increased by varieties compared to other methods. The use of mineral fertilizers in different ways in typical sierozem soil conditions affects the grain yield of local and foreign varieties. In other words, the average yield of medium-ripe soybean varieties "Nafis" was 43.4 c / ha, "Vilana" was 42.4 c / ha, and the best way to increase the yield is to apply fertilizers as NPK in combination with liquid fertilizer [17]. According to data of Khayrullayev Sardor Shamsiddin o'g'li and Usmonova Shohista Usmon qizi, the location of the lower first pod in soybean varieties is 12.8-15.9 cm in Orzu variety, 3-3.1 cm in Radimax stimulator, 2.2-2.4 cm in Gummat stimulator, 2.1 cm in Tecamin stimulator and 3.1 cm in Algora stimulator was found to be high. The most effective results were observed in Radimax, Gummat and Algora bio-simulators, and the location of the lower first pod was detected 14.7-17.6 cm in the "Nafis" variety, which was 2.5-2.9 cm higher in the Radimax stimulator, 2.2-2.5 cm higher in the Gummat stimulator, 2.1 cm higher in the Tecamine stimulator, and 2.4 cm higher in the Algora stimulator than in the control variant. The most effective results were observed in Radimax, Gummat and Algora biosimulators [15]. According to Atabayeva, K.N., Umarova, N.S., Yakubov, S., & Khayrullaev, S.S, positive results were obtained from moderate levels of sulphur and manganese, and low levels of iron. Macro and micronutrients had a positive effect on soy yield. An additional 7.6 quintals (q)/ha was harvested in exchange for macro fertilizer. Compared to the background variant, the yield was 4.6-8.3 q/ha for sulphur and 4.9-9.8 q/ha for manganese. The yield of the iron element was lower than that of the background variant. Grain quality has changed in exchange for macro and micronutrients. In exchange for mineral fertilizers, this figure increased by 2.4%. In exchange for the element sulphur, the protein increased by 3.1-5.8%; an increase of 4.4-8.4% was observed in exchange for the element manganese. It was noted that the protein increased by 7.9-8.7% in exchange for the element iron [18]. According to Ugli Khayrullayev, S.S., & Kizi Usmonova, S.U., mineral fertilizers and sulfur microelements activate the symbiotic activity of soybean variety "Orzu", averaging 32.4-42.3 million nodules per hectare, the number of nodules due to the background of mineral fertilizers increased by 13.6%, and there was an increase of 19.4-23,4% due to sulfur, as well as an average weight of nodules was 6.46-9.56 c / ha, the weight of nodules increased by 5.3% due to mineral fertilizers, and 17.1-32.4% due to sulfur. During the application period, 6.46-9.56 c / ha of nodules mass was accumulated per hectare according to the studied variants, which contributes to the increase of nitrogen and organic matter in the soil and a slight increase in biological efficiency [19]. According to Usmonova Sh.U, Khayrullaev S.Sh, Shomuqimov N.N, & Gaynanova A.F, the influence of stimulants on soybeans affected the weight of 1000 grains of Vilana cultivar, under the influence of Gummat stimulator this figure was 2.2-7.4 grams higher than on basis of mineral fertilizers (Background), and under the influence of Rival stimulator-3.0-6.0 grams [20]. According to Khayrullaev S.S, In the variant, where not used mineral fertilizers and micronutrients, the leaf area in the control variant of the Orzu variety of soybean was 51.1

thousand m<sup>2</sup> / ha. Under the influence of microelements, the leaf area of Orzu was 59.1-64.6 thousand m<sup>2</sup> / ha. The highest rates of exposure to micronutrients were observed with medium use of sulfur and manganese. Under influence macro and micro fertilizers, the leaf area of Orzu variety increased from 4.0 to 13.5 thousand m<sup>2</sup> / ha, or from 7.3 to 20.9% [21].

### **METHODS AND MATERIALS**

Experiments are carried out in field and laboratory conditions. In the research "Methods of conducting field experiments" (T.UzPITI 2007), "Methodology of field experiment (B.Dospekhov, 1985), "Methodology of the State variety testing of agricultural crops" (1985, 1989), "Methods of agrochemical, agrophysical studies of the soil of Central Asia" (1988) methods are used.

**Place and conditions of experiment.** The experiments were conducted in the experimental fields of Tashkent State Agrarian University. In this case, the experimental variants were in 4 repetitions for each variety, the number of variants was 10, the number of plots was 40, the length of each plot was 10 m, the width was 2.8 m. The variants were placed in a randomized method. Each option had 4 rows, and the total area of each plot was 28.0 m<sup>2</sup>, of which 2 rows in the middle were designated as counting rows, and 2 rows on the edge were designated as protection rows. The number of counting plants was 20. The experiments were carried out in field and laboratory conditions.

**Nafis variety.** The variety was created by the method of individual selection at the Rice Research Institute of Uzbekistan.

The growing period is 115-120 days. The height of the plant is 145-150 cm. The location of the lower pod is 14-16 cm, the number of branches is 2-4, the number of pods in one plant is 120-130, the number of grains in one pod is 2-4.

Grain quality and technological parameters: weight of 1000 seeds is 165-175 g. The protein content of grain is 40-41%, the oil content is 25-27%. Resistant to lodging, shedding, diseases and mechanized harvesting.

Yield: 30-32 q/ha grain yield and 250-300 q/ha blue mass can be obtained from the variety under favorable conditions.

**Vilana varieties:** The variety was created at the All-Russian Research Institute of Oil crops. This variety was obtained by cross-breeding the L-309 hybrid generation with the 0240 collection sample and individual and mass selection in the F<sub>2</sub> and F<sub>3</sub> varieties. The plant hairs are gray. The flowers are purple, the pods are brown, the seeds are yellow, dull, without spots. The variety is mid-ripening, resistant to adverse conditions, drought-resistant, and yields increase when watered. The growing season is 116-120 days. Seed yield without irrigation is 32-34 q/ha, and with irrigation up to 42 q/ha. Plant height is 111-115 cm, the lower pod location is 16-17 cm. The protein content in the grain is 40.1-40.3%, the oil content is 22.4-22.6%.

### **RESULTS AND DISCUSSION**

The effect of stimulants also has a positive effect on the location of the first pod, as a result of which the crop can be harvested without loss. In 2023, the height of the first pod in the Nafis variety when applied without fertilizer was 14.2 cm; when applied with mineral fertilizers, it was 14.9 cm, and when applied with mineral fertilizers in combination with water, it was 15.3 cm. When using the

stimulants Uzbiogumin, Regoplant, and Maltamin, the Maltamin stimulator showed better results than other stimulants, increasing by 2.7 cm compared to the case without fertilizer, by 2.0 cm compared to the case with mineral fertilizers, and by 1.6 cm compared to the case with mineral fertilizers in combination with water. When using complex silicon fertilizer, the height of the first pod location increased by 3.2 cm compared to the case without fertilizer, by 2.5 cm compared to the case with mineral fertilizers, and by 2.1 cm compared to the case with mineral fertilizers combined with water. When using complex silicon fertilizer in combination with Uzbiogumin, Regoplant, and Maltamin stimulants, the height of the first pod location increased by 4.1 cm compared to the case without fertilizer, by 3.4 cm compared to the case with mineral fertilizers, and by 3.0 cm compared to the case with mineral fertilizers combined with water.

In 2024, the height of the first pod location in the Nafis variety when used without fertilizer was 14.0 cm; When using mineral fertilizers, it was 14.7 cm and when using mineral fertilizers together with water, it was 15.1 cm. When using the stimulants Uzbiogumin, Regoplant and Maltamin, the Maltamin stimulator showed better results than other stimulants, increasing by 2.6 cm compared to the case without fertilizer, by 1.9 cm compared to the case with mineral fertilizers, and by 1.2 cm compared to the case with mineral fertilizers combined with water. When using a silicon complex fertilizer, the height of the first pod location increased by 3.1 cm compared to the case without fertilizer, by 2.4 cm compared to the case with mineral fertilizers, and by 2.0 cm compared to the case with mineral fertilizers combined with water. When using the complex silicon fertilizer in combination with Uzbiogumin, Regoplant and Maltamin stimulants, the height of the first pod was recorded as the highest when used with the Maltamin stimulator, increasing by 4.0 cm compared to the case without fertilizer, by 3.3 cm compared to the case with mineral fertilizers, and by 2.9 cm compared to the case with mineral fertilizers in combination with water.

In 2025, the height of the first pod was 14.9 cm when used without fertilizer in the Nafis variety; When using mineral fertilizers, it was 15.6 cm and when using mineral fertilizers together with water, it was 16.1 cm. When using the stimulants Uzbiogumin, Regoplant and Maltamin, the Maltamin stimulator showed better results than other stimulants, increasing by 2.8 cm compared to the case without fertilizer, by 2.1 cm compared to the case with mineral fertilizers, and by 1.6 cm compared to the case with mineral fertilizers combined with water. When using a complex silicon fertilizer, the height of the first pod increased by 3.4 cm compared to the case without fertilizer, by 2.7 cm compared to the case with mineral fertilizers, and by 2.2 cm compared to the case with mineral fertilizers combined with water. When using the silicon complex fertilizer in combination with the Uzbiogumin, Regoplant, and Maltamin stimulants, the height of the first pod was the highest when used with the Maltamin stimulator, increasing by 4.3 cm compared to the case without fertilizer, by 3.6 cm compared to the case with mineral fertilizers, and by 3.1 cm compared to the case with mineral fertilizers and water.

**Table 1**  
**The effect of stimulants and complex silicon fertilizers on the formation of lower pods of soybean varieties, (2023-2025)**

Rates for the use of stimulants and complex silicon fertilizers	Lower pods placement			
	2023	2024	2025	averag

				e
<b>Nafis</b>				
Control (without fertilizer)	14.2	14.0	14.9	14.4
Background - (N-60, P-120, K-80 kg/ha)	14.9	14.7	15.6	15.1
Background - (N-60, P-120, K-80 kg/ha)+suv	15.3	15.1	16.1	15.5
Background +Uzbiogumin (1,8 l/ha)	15.9	15.7	16.7	16.1
Background +Regoplant (150 ml/ha)	16.2	16.0	17.0	16.4
Background +Maltamin (3,5 l/ha)	16.9	16.6	17.7	17.1
Background +Sila kremniya (450 ml/ha)	17.4	17.1	18.3	17.6
Background +Sila kremniya+ Uzbiogumin (450 ml/ha+1,8 l/ha)	17.7	17.4	18.6	17.9
Background +Sila kremniya+ Regoplant (450 ml/ha+150 ml/ha)	18.0	17.7	18.9	18.2
Background +Sila kremniya+ Maltamin (450 ml/ha+3,5 l/ha)	18.3	18.0	19.2	18.5
<b>Vilana</b>				
Control (without fertilizer)	12.5	12.3	13.1	12.6
Background - (N-60, P-120, K-80 kg/ha)	13.1	12.9	13.8	13.3
Background - (N-60, P-120, K-80 kg/ha)+suv	13.5	13.3	14.1	13.6
Background +Uzbiogumin (1,8 l/ha)	14.0	13.8	14.7	14.2
Background +Regoplant (150 ml/ha)	14.3	14.0	15.0	14.4
Background +Maltamin (3,5 l/ha)	14.9	14.6	15.6	15.0
Background +Sila kremniya (450 ml/ha)	15.3	15.1	16.1	15.5
Background +Sila kremniya+ Uzbiogumin (450 ml/ha+1,8 l/ha)	15.6	15.3	16.4	15.8
Background +Sila kremniya+ Regoplant (450 ml/ha+150 ml/ha)	15.8	15.6	16.6	16.0

Background +Sila kremniya+ Maltamin (450 ml/ha+3,5 l/ha)	16.1	15.9	16.9	16.3
---	------	------	------	------

Based on average 3-year data, the height of the first pod in the Nafis variety when applied without fertilizer was 14.4 cm; when applied with mineral fertilizers, it was 15.1 cm; and when applied with mineral fertilizers in combination with water, it was 15.5 cm. When using the stimulants Uzbiogumin, Regoplant, and Maltamin, the Maltamin stimulant showed better results than other stimulants, increasing by 2.7 cm or 18.8% compared to the one applied without fertilizer, by 2.0 cm or 13.2% compared to the one applied with mineral fertilizers, and by 1.6 cm or 10.3% compared to the one applied with mineral fertilizers in combination with water. When using complex silicon fertilizer, the height of the first ear of corn increased by 3.2 cm or 22.2% compared to the case without fertilizer, by 2.5 cm or 16.6% compared to the case with mineral fertilizers, and by 2.1 cm or 13.5% compared to the case with mineral fertilizers and water. When using complex silicon fertilizer in combination with Uzbiogumin, Regoplant and Maltamin stimulants, the height of the first ear of corn increased by 4.1 cm or 28.5% compared to the case without fertilizer, by 3.4 cm or 22.5% compared to the case with mineral fertilizers, and by 3.0 cm or 19.4% compared to the case with mineral fertilizers and water.

In the Vilana variety, the height of the first pod in 2023 was 12.5 cm without fertilizer; 13.1 cm with mineral fertilizers and 13.5 cm with mineral fertilizers in combination with water. When using Uzbiogumin, Regoplant and Maltamin stimulants, the Maltamin stimulant showed better results than other stimulants, increasing by 2.4 cm compared to the case without fertilizer, by 1.5 cm compared to the case with mineral fertilizers and by 1.4 cm compared to the case with mineral fertilizers in combination with water. When using complex silicon fertilizer, the height of the first pod location increased by 2.8 cm compared to the case without fertilizer, by 2.2 cm compared to the case with mineral fertilizers, and by 1.8 cm compared to the case with mineral fertilizers in combination with water. When using complex silicon fertilizer in combination with Uzbiogumin, Regoplant, and Maltamin stimulants, the height of the first pod location increased by 3.6 cm compared to the case without fertilizer, by 3.0 cm compared to the case with mineral fertilizers, and by 2.6 cm compared to the case with mineral fertilizers in combination with water.

In 2024, the height of the first pod location of Vilana was 12.3 cm when used without fertilizer; When using mineral fertilizers, it was 12.9 cm and when using mineral fertilizers together with water, it was 13.3 cm. When using the stimulants Uzbiogumin, Regoplant and Maltamin, the Maltamin stimulant showed better results than other stimulants, increasing by 2.3 cm compared to the case without fertilizers, by 1.7 cm compared to the case with mineral fertilizers and by 1.3 cm compared to the case with mineral fertilizers and water. When using complex silicon fertilizer, the height of the first pod increased by 2.8 cm compared to the case without fertilizers, by 2.2 cm compared to the case with mineral fertilizers and by 1.8 cm compared to the case with mineral fertilizers and water. When using the complex silicon fertilizer in combination with Uzbiogumin, Regoplant and Maltamin stimulants, the height of the first pod was recorded as the highest when used with the Maltamin stimulant, increasing by 3.6 cm compared to the case without fertilizer, by 3.0 cm compared to the case with mineral fertilizers, and by 2.6 cm compared to the case with mineral fertilizers and water.

In 2025, the height of the first pod was 13.1 cm in the case without fertilizer; When using mineral fertilizers, it was 13.8 cm and when using mineral fertilizers together with water, it was 14.1 cm. When using Uzbiogumin, Regoplant and Maltamin stimulants, the Maltamin stimulant showed better results than other stimulants, increasing by 2.5 cm compared to the case without fertilizers, by 1.8 cm compared to the case with mineral fertilizers, and by 1.5 cm compared to the case with mineral fertilizers combined with water. When using silicon complex fertilizer, the height of the first pod increased by 3.0 cm compared to the case without fertilizers, by 2.3 cm compared to the case with mineral fertilizers, and by 2.0 cm compared to the case with mineral fertilizers combined with water. When using the silicon complex fertilizer in combination with the stimulants Uzbiogumin, Regoplant, and Maltamin, the height of the first pod was the highest when used with the Maltamin stimulant, increasing by 3.8 cm compared to the case without fertilizer, by 3.1 cm compared to the case with mineral fertilizers, and by 2.8 cm compared to the case with mineral fertilizers and water.

Based on the average 3-year data, the height of the first pod in the Vilana variety was 12.6 cm when applied without fertilizer; 13.3 cm when applied with mineral fertilizers, and 13.6 cm when applied with mineral fertilizers in combination with water. When using the stimulants Uzbiogumin, Regoplant, and Maltamin, the Maltamin stimulant showed better results than other stimulants, increasing by 2.4 cm or 19.0% compared to the one applied without fertilizer, by 1.7 cm or 12.8% compared to the one applied with mineral fertilizers, and by 1.4 cm or 10.3% compared to the one applied with mineral fertilizers in combination with water. When using complex silicon fertilizer, the height of the first ear of corn increased by 2.9 cm or 23.0% compared to the case without fertilizer, by 2.2 cm or 16.5% compared to the case with mineral fertilizers, and by 1.9 cm or 14.0% compared to the case with mineral fertilizers combined with water. When using complex silicon fertilizer in combination with Uzbiogumin, Regoplant, and Maltamin stimulants, the height of the first ear of corn increased by 3.7 cm or 29.4% compared to the case without fertilizer, by 3.0 cm or 22.6% compared to the case with mineral fertilizers, and by 2.7 cm or 19.9% compared to the case with mineral fertilizers combined with water.

Good results were observed when the Maltamin (3.5 l/ha) stimulator, Sila silicon fertilizer (450 ml/ha) and Fon + Sila silicon + Maltamin (450 ml/ha+3.5 l/ha) were applied to the height of the first pod.

## **CONCLUSIONS**

Therefore, it is known from the experimental results that stimulants and complex fertilizers with silicon had a positive effect on the height of the first pod of soybean varieties, and in both varieties, when the stimulants were Maltamin stimulator, when sila silicon was compared with stimulants, when sila silicon was used together with stimulants, and when sila silicon+Maltamin was used together, the height of the first pod was higher than in other options.

**REFERENCES**

1. Агафонов О.М. и др. Научные основы применения удобрений. Колос. 2015. с 11,187.
2. Доспехов Б."Методика полевого опыта, 1985 й
3. Балакай Г.Т., Ивевбор Лоуренс Уче и др. Масличные культуры, 2008
4. Ларина Р.Е., Лисова Р.В., Логинов О.Н., Ловцова. Особенности формирования урожая сои в условиях центральной зоны Нечерноземной зоны // Проблемы агрохимии и экологии. 2018, № 4,С.27-33
5. Ран.О.П.,Селихова.О.А.,Тихончук.П.В. Применение биологических препаратов в посевах сои // Достижения науки и техники АПК. 2009, №18.С.26-27
6. Сонин К.Е. Влияние препарата фуrolан на формирование качества семян трёх сортов подсолнечника // Пищевая технология. – 2010. – №1. – С. 13–15.
7. Шаповал, Новицкая.Н.В., Щучка Р.В, Джелисюк.А.В. Урожайность сои в зависимости от элементов технологии на черноземах типичных лесостепи Украины // Вестник. Алтайский ГАУ, №5. серия 11. 2015
8. Хохоева Н.Т., Ран.О.П. Применение биологических препаратов в посевах сои // Достижения науки и техники АПК. 2009, №18.С.26-27.
9. Якубжонов.О, С.Турсунов, Мукимов.З Дончилик Т. Янги аср авлоди 2009 234-бет
10. Nazarovna, AK, Bakhromovich, NF, Alavkhonovich KA, Ugli KSS. Effects of Sulfur and Manganese Micronutrients on the Yield of Soybean Varieties. *Agricultural Sciences*, 2020;11: 1048-1059. <https://doi.org/10.4236/as.2020.1111068>
11. JURAEVA R., TASHPULATOV J., IMINOV A., BOZOROV X., ZAYNITDINOVA L., & KUKANOVA S. (2020). EFFICIENCY OF SYMBIOTIC NITROGEN FIXATION OF SOY NODULE BACTERIA AFTER PRESERVATION. *PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY*, 21(61-62), 7279. Retrieved from <https://www.ikpress.org/index.php/PCBMB/article/view/5644>.
12. ABDUMANNOBOVICH, I. A., UGLI, K. S. S., ERGASHBOYEVNA, K. D., UGLI, H. S. R., & TURSINALIEVICH, K. S. (2021). EFFECTS OF NITRAGINE AND MINERAL FERTILIZER RATES USED ON REPEATED LEGUME CROPS ON SEED GERMINABILITY. *PLANT CELL BIOTECHNOLOGY AND MALECULAR BIOLOGY*, 22(35-36), 251-258. Retrieved from <https://www.ikpress.org/index.php/PCBMB/article/view/6342>.
13. Atabayeva Khalima Nazarovna1, Khayrullaev Sardor Shamsiddin o'g'li, and Usmonova Shohista Usmon qizi, "THE EFFECT OF SULPHUR ON THE BRANCHING OF SOYBEAN VARIETIES", *STJITT*, vol. 1, no. 2, pp. 4-6, Aug. 2020.
14. Iminov, A. A., Hatamov, S. R. O., & Khayrullaev, S. S. O. (2020). Effect Of Nitragine And Mineral Fertilizers On Soil Microbiological Properties In Planted As Secondary Legume Crops. *The American Journal of Agriculture and Biomedical Engineering*, 2(08), 169-172. <https://doi.org/10.37547/tajabe/Volume02Issue08-22>
15. Khayrullayev Sardor Shamsiddin o'g'li and Usmonova Shhista Usmon qizi, "THE EFFECT OF STIMULATORS ON THE LOCATION OF THE FIRST POD OF SOYBEAN VARIETIES", *STJITT*, vol. 1, no. 2, pp. 7-9, Aug. 2020.

16. Khayrullayev Sardor Shamsiddinugli. Effect Of Sulfur And Manganese Micro Nutrients On Grain Quality Of Soybean Varieties. *ACADEMICIA: An International Multidisciplinary Research Journal* <https://saarj.com>. Vol. 11, Issue 3, March 2021. DOI: 10.5958/2249-7137.2021.00582.6. P.63-68.

17. Umarova Nigora Sadridinovna, Bo'riboev Bekzod Yetmish ugli, Khayrullayev Sardor Shamsiddin ugli, Usmonova Shokhista Usmon kizi, & Turdaliyeva Shokhista Tulkinjon kizi. (2021). EFFECT OF MINERAL FERTILIZERS ON MORPHOPHYSIOLOGICAL CHARACTERISTICS OF LOCAL AND FOREIGN SOYBEAN VARIETIES. *European Journal of Agricultural and Rural Education*, 2(4), 39-42. Retrieved from <https://scholarzest.com/index.php/ejare/article/view/611>.

18. Atabayeva, K. N., Umarova, N. S., Yakubov, S., & Khayrullaev, S. S. (2021, December). Influence of trace elements on soybean yield in grassland-swamp soils. In *IOP Conference Series: Earth and Environmental Science* (Vol. 939, No. 1, p. 012049). IOP Publishing.

19. Ugli Khayrullayev, S. S., & Kizi Usmonova, S. U. (2021). Effect Of Sulfur Micronutrient On Symbiotic Activity Of “Orzu” Variety Of Soybean. *The American Journal of Agriculture and Biomedical Engineering*, 3(05), 49–59. <https://doi.org/10.37547/tajabe/Volume03Issue05-10>

20. Usmonova Sh.U, Khayrullaev S.Sh, Shomuqimov N.N, & Gaynanova A.F. (2022). Effect Of Stimulators On Thousand Piece Grain Weight Of “Vilana” Cultivar Of Soybean. *Central Asian Journal of Theoretical and Applied Science*, 3(4), 153-157. Retrieved from <https://cajotas.centralasianstudies.org/index.php/CAJOTAS/article/view/467>

21. Khayrullaev S. S. Influence of sulfur and manganese micronutrients on formation of leaf area of “Orzu” cultivar of soybean // *ISJ Theoretical & Applied Science*, 03 (107). – 2022. – C. 287-191.